


Constraints on smoke injection height, source strength, and transports from MISR and MODIS

Ralph Kahn


NASA Goddard Space Flight Center

Mariya Petrenko, Maria Val Martin, Mian Chin



The NASA Earth Observing System's Terra Satellite

*First Light:
February 24, 2000*



Source: Terra Project Office / NASA Goddard Space Flight Center


Multi-angle Imaging SpectroRadiometer



<http://www-misr.jpl.nasa.gov>
<http://eosweb.larc.nasa.gov>

- Nine CCD push-broom cameras
- Nine view angles at Earth surface:
70.5° forward to 70.5° aft
- Four spectral bands at each angle:
446, 558, 672, 866 nm
- *Studies Aerosols, Clouds, & Surface*


Ten Years of Seasonally Averaged Mid-visible Aerosol Optical Depth from **MISR**



...includes bright desert dust source regions



MISR *Aerosol Type* Distribution

MISR Version 22, July 2007




Wildfire Smoke Injection Heights & Source Strengths


[These are *the two key parameters* representing aerosol sources in climate models]



Val Martin et al. ACP 2010



MODIS Smoke Plume Image & Aerosol Amount Snapshots




Different Techniques for Assuming Model Source Strength
Overestimate or **Underestimate** Observation
Systematically in Different Regions


GoCART Model-Simulated Aerosol Amount Snapshots
for Different Assumed Source Strengths

Petrenko et al., JGR 2012

Changes in geometric perspective with angle






Changes in geometric perspective with angle










MISR *Smoke Plume Height* Mapping

July 2, 2004, Canada near Alaska border




Oregon Fire Sept 04 2003

Orbit 19753 Blks 53-55 MISR Aerosols V17, Heights V13 (no winds)




Detail of Wildfire Source Region


Oregon Fire Sept 04 2003




MISR Nadir **275 m** Image



MODIS Image + **Fire Power**




MISR **Plume Heights** for Sub-patches



Very Simple Plume Parcel Model

→ **Broad swath + high spatial resolution** needed to characterize sources


N. America Plume *Injection Height* Climatology



~ 3400 plumes digitized over North America for 2002, 2004-2007

- Tropical Forest
- Temperate Forest
- Boreal Forest
- Boreal Shrubland
- Non-Boreal Shrubland
- Boreal Grassland
- Non-Boreal Grassland
- Cropland


MODIS IGBP land cover map
(1x1 Km res)



Percent of plumes >0.5 km *above BL*, stratified by year and vegetation type

Evaluation of a 1D plume-rise model: Towards a parameterization of smoke *injection heights*


To Constrain models:
Need to assess the
Parameterizations
actually used



1-D Plume-rise model heights vs. MISR-observed max. plume heights

-- Plume-rise calculations have *lower dynamic range than observed*, but very variable

Evaluation of a 1D plume-rise model: Towards a parameterization of smoke *injection heights*




The key factors:


- *Fire Energy*
(fire area; heat flux, FRP)
- *Atmospheric Stability*
- *Entrainment*

Plume height increases systematically as *FRP* increases and *Atmospheric Stability* decreases

Satellite AOD snapshots to constrain Biomass Burning Emissions *Source Strength*






- 1 Tree cover, broadleaved, evergreen
- 2 Tree cover, broadleaved, deciduous, closed
- 3 Tree cover, broadleaved, open
- 4 Tree cover, needle-leaved, evergreen
- 5 Tree cover, needle-leaved, deciduous
- 6 Tree cover mixed leaf type
- 7 Tree cover, regularly flooded, fresh water
- 8 Tree cover, regularly flooded, saline water
- 9 Mosaic: tree cover / other natural vegetation
- 10 Undefined
- 11 Shrub cover, closed-open, evergreen
- 12 Shrub cover, closed-open, deciduous
- 13 Herbaceous cover, closed-open
- 14 Sparse herbaceous or sparse shrub cover
- 15 Regularly flooded shrub and/or herbaceous cover
- 16 Cultivated and managed areas
- 17 Mosaic: Cropland/Tree cover/other natural veg
- 18 Cropland/Shrub and/or grass cover



13 Smoke Emission Estimates

MODIS-GoCART Total Column AOD Comparisons

Sample Case: Siberia July 20 2006



**Goddard Chemistry Aerosol
Radiation and Transport (GOCART)
model runs**

3-hourly output













Resolution: 1°(lat) x 1.25°(lon) x 30 vert.
layers

Meteorological fields GEOS DAS Version 4

Emissions include: dust, sea salt,
anthropogenic, sulfate & precursors, BB

13 BB emission options in separate model
runs

Study period: June 2006-June 2007




AOD (550nm)

Petrenko et al., JGR, 2012

Ratio of GOCART to MODIS average AOD

For each case, for 12 emission estimates



Systematic regional patterns; some emissions work better in certain regions

Ratio of GOCART average AOD to MODIS average AOD




Quantitative Relationship Between Smoke Emission and AOD


Depends On

- *Wind Speed at source*
- *Background AOD*

Steeper slope ~
Lower wind speed





High background AOD ~
Smoke plume insignificant



SEAC4RS Field Campaign

DC-8 and ER-2 Flights Monday, 19 August 2013



MISR (Multi-angle Imaging SpectroRadiometer) Overpass




Monday, 19 August 2013 17:40 UTC

17:40 UTC
Path 031
Orbit 72716

South Dakota

Nebraska



Kansas



MISR Aerosol Optical Depth (Research Algorithm)

19 August 2013


Site 2



MISR Aerosol Type (Research Algorithm)

19 August 2013

Site 2



Smoke Plume 1

AOD 0.35-0.9

ANG 1.5-1.9 (*small*)

SSA 0.94-0.98 (*absorbing*)

FrNon-Sph 0-0.2 (*mostly spherical*)

Continental Background

AOD 0.15-0.2

ANG 1.0-1.5 (*medium*)

SSA 0.99-1.0 (*non-absorbing*)

FrNon-Sph 0.0 (*spherical*)

Smoke Plume 2

AOD 0.35-0.6

ANG 1.6-2.0 (*smaller*)

SSA 0.96-0.98 (*less absorbing*)




FrNon-Sph 0-0.1 (*more spherical*)

70° Forward View

Passive-remote-sensing **Aerosol Type** is a **Total-Column-Effective, Categorical** variable!!

Site 2 Smoke Transports


19 August 2013



DIAL – Hair et al.





GEOS-5 MODEL Aerosol Optical Depth

19 August 2013 18 UTC




GEOS-5 MODEL Aerosol Type

19 August 2013 18 UTC



AeroCom BB Experiment AOD – *Motivation*



- We have a substantial set of **satellite wildfire plume AOD snapshots and injection heights** to help calibrate model/inventory performance
- We are: (1) adding **more fire source-strength cases**, (2) using MISR to **improve the AOD constraints** and (3) adding **2008 global injection heights**
- We selected **GFED3-daily** due to good overall source strength performance, but **any inventory can be tested**
- Joint effort, to **test multiple, global models** to draw robust BB injection height & emission strength conclusions

We provide: Satellite-based **injection height** and **smoke plume AOD** climatologies

Experiment Design

	Exp.	BB Daily emission	Injection height
	BB0	<i>No BB emission</i>	
Control →	BB1	GFED v3	Boundary layer
	BB2	GFED v3 x 0.5	Boundary layer
	BB3	GFED v3 x 2	Boundary layer
	BB4	GFED v3 x 5	Boundary layer
Stage 2 →	BB5	GFED v3	From MISR plume ht.
	BB6	GFED v3 x 5	From MISR plume ht.

Requested output:

2-D, 3-hourly, instantaneous


- Total column **550 nm AOD**
- **Biomass burning AOD**, if available (or AOD's of individual aerosol species)
- **Wind speeds** in the middle of emission injection height
[e.g., if all smoke is distributed within PBL, output mid-PBL winds]
- **PBL height**

3-D [3-hourly]


- Aerosol **species concentrations**
- Aerosol 550 nm **extinction**

With *Source Strength* Perturbation Factors: 0.7, 1, 3 & 5


GOCART GFED3x0.7




GOCART GFED3




GOCART GFED3x3



GOCART GFED3x5



Ratio of GOCART average AOD to MODIS average AOD





Satellites

frequent, global
snapshots;
aerosol amount &
aerosol type maps,
plume & layer heights

**Aerosol-type
Predictions**

Model Validation

- Parameterizations
- Climate Sensitivity
- Underlying mechanisms

Remote-sensing Analysis

- Retrieval Validation
- Assumption Refinement

Regional Context

CURRENT STATE

- Initial Conditions
- Assimilation


Suborbital



targeted chemical &
microphysical detail



point-location
time series



Models

space-time interpolation,

**DARF &
Anthropogenic
Component**

calculation and prediction